

# Physics 681: Solar Physics and Instrumentation – Lecture 15

*Carsten Denker*

*NJIT* Physics Department  
Center for Solar–Terrestrial Research



## Convection

- Stability
- Mixing-Length Theory
  - The Local Formalism
  - Numerical Test Calculations
  - Overshooting: A Non-Local Formalism
- Granulation
- Mesogranulation
- Supergranulation
- Giant Cells

October 27, 2005

Center for Solar-Terrestrial Research



# Stability

- Energy transport mechanisms: radiation, convection, and conduction
- Stratification of the outer layer ( $\approx 200,000$  km) is unstable
- Mixing-length concept: parcel of gas travels a certain distance, dissolves, and deposits its energy
- Adiabatic displacement from equilibrium position (no oscillation!)
- Schwarzschild criterion (1906)
- Condition for instability (pressure equilibrium)

$$\rho^* - \rho_0^* = \delta r \left[ \left( \frac{d\rho}{dr} \right)_a - \frac{d\rho}{dr} \right] < 0$$

- Variations of the mean molecular weight  $\mu$  and chemical composition

$$\frac{dT}{dr} < \left( \frac{dT}{dr} \right)_a + \frac{T}{\mu} \frac{d\mu}{dr} - \frac{T}{\mu} \left( \frac{d\mu}{dr} \right)_a$$



October 27, 2005

Center for Solar-Terrestrial Research

- Ideal gas equation  
 $P = \rho \mathcal{R} T / \mu$
- Core material is fully ionized ( $d\mu/dr$ )<sub>a</sub> = 0
- $d\mu/dr$  towards center is negative due to the accumulation of helium  
⇒ stabilizing
- Assume instantaneous adjustment of the ionization equilibrium

$$\frac{d\mu}{dr} = \left( \frac{\partial \mu}{\partial P} \right)_T \frac{dP}{dr} + \left( \frac{\partial \mu}{\partial T} \right)_P \frac{dT}{dr} \text{ and}$$

$$\frac{d\mu}{dr} = \left( \frac{\partial \mu}{\partial P} \right)_T \left( \frac{dP}{dr} \right)_a + \left( \frac{\partial \mu}{\partial T} \right)_P \left( \frac{dT}{dr} \right)_a$$

- Schwarzschild criterion

$$\frac{dT}{dr} < \left( \frac{dT}{dr} \right)_a$$

$$\nabla > \nabla_a \Leftrightarrow \frac{d \ln T}{d \ln P} > \left( \frac{d \ln T}{d \ln P} \right)_S$$



October 27, 2005

Center for Solar-Terrestrial Research

# Mixing-Length Theory

- Viscosity vs. turbulent motion
- Numerical methods (Navier-Stokes equation in 3D  $\mathcal{N}$ )
- Schmidt and Prandtl (1915-1930) and Böhm-Vitense (1958)
- Mixing-length

$$l = \alpha H_p$$

- Energy flux

$$F_R + F_C = L_\odot / 4\pi r^2$$

- Radiative flux

$$F_R = -\frac{16\sigma T^3}{3\kappa\rho} \frac{dT}{dr} = \frac{16\sigma T^4}{3\kappa\rho H_p} \nabla$$

- Mean excess temperature

$$\Delta T = \left[ \left( \frac{dT}{dr} \right)' - \frac{dT}{dr} \right] \delta r \equiv (\nabla - \nabla') \frac{T \delta r}{H_p}$$



October 27, 2005

Center for Solar-Terrestrial Research

- Assume that the mean value of  $\delta r$  is half the mixing length

$$\Delta T = (\nabla - \nabla') T \alpha / 2$$

- Convective flux

$$F_C = \alpha \rho c_p v T (\nabla - \nabla') / 2$$

- Convective velocity

$$\partial^2 \delta r / \partial t^2 = -g \Delta \rho / \rho = g \delta \Delta T / T$$

- Pressure equilibrium ( $\Delta P = 0$ ) and integration

$$\left( \frac{\partial \delta r}{\partial t} \right)^2 = \frac{g \delta}{H_p} (\nabla - \nabla') (\delta r)^2$$

- Work done by buoyancy force appears as kinetic energy of the parcel

- Introduce correction factor of  $1/2$  to account for friction

$$v = l \left[ \frac{g \delta}{8 H_p} (\nabla - \nabla') \right]^{1/2}$$



October 27, 2005

Center for Solar-Terrestrial Research

- Radiative losses across the surface of the parcel ( $d$  = distance over which  $\Delta T$  drops to zero)

$$F'_R = -\frac{16\sigma T^3}{3\kappa\rho} \frac{\Delta T}{d} = \frac{8\alpha\sigma T^4}{3\kappa\rho d} (\nabla' - \nabla)$$

- Convective flux

$$F_C = \alpha\rho c_p v T (\nabla - \nabla_a) / 2 + \alpha\rho c_p v T (\nabla_a - \nabla') / 2$$

- First term (ideal adiabatic conditions) and second term (real convective flux)

- Radiative loss per unit time

$$\frac{16\sigma T^3 S}{3\kappa\rho d} (\nabla' - \nabla) = \rho c_p v q (\nabla_a - \nabla')$$

- Spherical parcels  $d = (8/9) l$  and  $S/l / qd = 9/2$

- Temperature gradients

$$\nabla' - \nabla_a = 2U (\nabla - \nabla') \quad \text{with} \quad U = \frac{24\sqrt{2}\sigma T^3 P^{1/2}}{c_p \kappa g l \delta^{1/2} \rho^{5/2}}$$



October 27, 2005

Center for Solar-Terrestrial Research

## Conditions for Convection in Stars

- The opacity is large, implying that an unachievable temperature gradient would be necessary for radiative energy transport
- A region exists where ionization is occurring, causing a large specific heat and a low adiabatic temperature gradient
- The local gravitational acceleration is, as would be the case in distended stars, again leading to a low adiabatic temperature gradient
- The temperature dependence of the nuclear energy generation rate is large, causing a steep radiative flux gradient and a large temperature gradient (CNO cycle or triple alpha process)



October 27, 2005

Center for Solar-Terrestrial Research